Evaluation of *kalrn* transcript levels in zebrafish seizure model

Marcela M. Simões (IC), Patrícia G. Barbalho (PQ), Cláudia V. Maurer-Morelli (PQ).

Abstract

Temporal transcript profile of *kalrn* gene in zebrafish larvae brain after pentylenetetrazole (PTZ)-induced seizure.

Key words: zebrafish, *kalrn*, seizure.

Introduction

Kalirin (*kalrn*) regulates neuronal shape and spine growth playing an important role in plasticity [1]. Besides, it has been implicated in some neurological diseases [2]. Today, there is no study associating the *kalrn* gene and seizures. We have been using the zebrafish seizure model for genetic investigations because its advantages in genetics studies. The main aim of this study was to investigate the *kalrn* mRNA levels in the zebrafish pentylenetetrazole (PTZ)-seizure model.

Results and Discussion

Our results showed no statistical difference in the *kalrn* mRNA levels between seizure group (SG) and control group (CG) after 24 and 48 hours of PTZ-induced seizure (images 1 and 2). The mean $\pm$ SEM of *kalrn* mRNA comparing SG and CG groups were the following: CG24h 0.85 ± 0.14 ; SG24h 0.92 ± 0.1 ($p=0.5$) and CG48h 1.00 ± 0.03 ; SG48h 0.93 ± 0.03 ($p=0.1$).

Image 1. Relative quantification of *kalrn* mRNA in zebrafish larvae brain after 24 hours after pentylenetetrazole-evoked seizure.

Image 2. Relative quantification of *kalrn* mRNA in zebrafish larvae brain after 48 hours after pentylenetetrazole-evoked seizure.

The *kalrn* gene has been associated with human neurological disorders, such as schizophrenia, Alzheimer disease, Huntington's disease and ischemic stroke[2]; nevertheless, there is no information about this gene and epilepsy. Although no differences were found in *kalrn* transcripts at 24h and 48h after PTZ, we cannot conclude that *kalrn* does not play a role after seizures.

Conclusions

By investigating other time points of *kalrn* mRNA expression we may shed some light into the role of this gene and its relation with the mechanisms underlying plasticity in epilepsy.

Acknowledgement

I would like to greatly thank the patience and help of my supervisor Claudia Morelli, and also Patricia Barbalho and all members of zebrafish lab. Supported by: PIBIC-CNPq and BRAINNCEPID/FAPESP, BRAZIL

¹ Ma XM et al. Kalirin-7 is required for synaptic structure and function. *J Neurosci* **2008**, 28, 12368–12382.

² Penzes P, Remmers C. Kalirin Signaling: Implications for Synaptic Pathology. *Mol Neurobiol* **2012**, 45(1), 109–118.