Screening of Lipases for Enantioselective Hydrolysis of 1-butyryloxyarylphosphonates.

Leticia M. Sato*, Taisa N. Kitamura, Lucidio C. Fardelone, José Augusto R. Rodrigues e Paulo J. S. Moran.

Abstract

The screening of lipases for use in enantioselective hydrolysis of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** furnished the methodology for obtention of alpha-hydroxy phosphonates. These alpha-hydroxy phosphonates that can be used as building blocks for pharmaceuticals and agrochemicals compounds.

Key words:

alpha-hydroxy phosphonates, biocatalysis, enantioselective hydrolysis, lipase.

Introduction

The optically active phosphorus compounds are important classes in organic chemistry, due to their biological properties, such as enzyme inhibitors, antibiotics, antiviral agents, antitumor and agrochemicals¹.

The use of lipases for resolution of racemic secondary alcohols are a great strategy in organic synthesis and in this work, we report the preliminary results for screening of enantioselective hydrolyses of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** by using *Aspergillus niger* AP12, *Burkholderia cepacia*, *Burkholderia cepacia* PS30, *Candida antarctica*, *Candida cylindracea* Type VII, *Candida rugosa*, and *Pseudomonas fluorescens* lipases.

Results and Discussion

The alpha-hydroxy phosphonates ($\pm$)-**2a-b** were synthesized using benzaldehyde, 3-chlorobenzaldehyde, respectively, 1.1 equivalent of trimethyl phosphite and oxalic acid (catalyst), without solvent, at 80°C². The esterification reaction of alpha-hydroxy phosphonates ($\pm$)-**2a-b** furnished the compounds ($\pm$)-**3a-b** by using butyric anhydride and DMAP in dichloromethane, at room temperature, Image 1.

The preliminary results of enantioselective hydrolysis of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** by using screening of lipases were demonstrated in Chart 1.

Image 1. Enantioselective hydrolysis of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** mediated by lipases.

Chart 1. Enantioselective hydrolysis of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** mediated by lipases.^a

Subst.	Lipase	Product	Conv. (%)	ee (%)	E (%)
3a	<i>A. niger</i> AP12	-	-	-	-
3a	<i>B. cepacia</i>	-	-	-	-
3a	<i>B. cepacia</i> PS30	-	-	-	-
3a	<i>C. antarctica</i>	2a	20	-	-
3a	<i>C. cylindracea</i>	(<i>R</i>)- 2a	40	6.0	1.2
3a	<i>C. rugosa</i>	(<i>R</i>)- 2a	40	31.5	2.3
3a	<i>P. fluorescens</i>	2a	40	-	-
3b	<i>A. niger</i> AP12	-	-	-	-
3b	<i>B. cepacia</i>	-	-	-	-
3b	<i>B. cepacia</i> PS30	-	-	-	-
3b	<i>C. antarctica</i>	2b	20	-	-
3b	<i>C. cylindracea</i>	(<i>R</i>)- 2b	40	16.3	1.5
3b	<i>C. rugosa</i>	(<i>R</i>)- 2b	40	69.2	8.6
3b	<i>P. fluorescens</i>	2b	40	-	-

^a 0.6 mmol of substrate dissolved in 3 mL of diisopropyl ether was added in 3 mL of phosphate buffer 0.1M, pH 7, containing 100 mg of Lipase, and stirred for 12 h at room temperature. The products were purified in chromatographic system Biotage using hexane/acetone gradient, and furnished isolated yields >90%. The configuration of alpha-hydroxy phosphonates was determined by comparison of the $[\alpha]_D^{20}$ with literature³ ((*R*)-**2a** $[\alpha]_D^{20}$ = +44.7, C 1.0 CHCl₃ and, (*R*)-**2b** $[\alpha]_D^{20}$ = +21.5, C 2.0, CHCl₃) and the enantiomeric excess was performed using HPLC by chiral column.

Conclusions

Enantioselective hydrolysis of 1-butyryloxyarylphosphonates ($\pm$)-**3a-b** mediated by lipases from *Candida cylindracea* and *Candida rugosa* furnished the alpha-hydroxy alcohols in 40% of conversion, and the studies are being conducted to optimize the biocatalytic process.

Acknowledgement

The authors are grateful for financial support of this project by the FAPESP (2014/00108-9), CAPES and CNPq.

¹ (a) Smirnov, I.; Belogurov Jr., A.; Friboulet, A.; Masson, P.; Gabibov, A.; Renard, P.-Y. *Chem.-Biol. Interact.* **2013**, *203*, 196. (b) Sobhani, S.; Tashrif, Z.; *Tetrahedron* **2010**, *66*, 1429. (c) Grembecka, J.; Mucha, A.; Cierpicki, T.; Kafarski, P., *J. Med. Chem.* **2003**, *46*, 2641.

² Vahdat, S. M.; Baharfar, R.; Tajbakhsh, M.; Heydari, A.; Bagbanian, S. M.; Khaksar, S., *Tetrahedron Lett.* **2008**, *49*, 6501.

³ Sun, L.; Guo, Q.-P.; Li, X.; Zhang, L.; Li, Y.-Y.; Da, C.-S. *Asian J. Org. Chem.* **2013**, *2*, 1031.